

Department of Health

IMMUNISATION SCHEDULE FOR CHILDREN

PROFMED



AGE OF CHILD	VACCINES NEEDED	HOW IT'S ADMINISTERED
AT BIRTH	BCG Bacillus Calmette Guerin	Right arm
	OPV (0) Oral Polio Vaccine	By mouth
6 WEEKS	OPV (1) Oral Polio Vaccine	By mouth
	RV (1) Rotavirus Vaccine	By mouth
	DTaP-IPV // Hib (1) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine and Haemophilus Influenzae Type b Combined	Intramuscular / Left thigh
	Hep B (1) Hepatitis B Vaccine	Intramuscular / Right thigh
	PCV ₇ (1) Pneumococcal Conjugated Vaccine	Intramuscular / Right thigh
10 WEEKS	DTaP-IPV // Hib (2) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine and Haemophilus Influenzae Type b Combined	Intramuscular / Left thigh
	Hep B (2) Hepatitis B Vaccine	Intramuscular / Right thigh
14 WEEKS	RV (1) Rotavirus Vaccine*	By mouth
	DTaP-IPV // Hib (3) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine and Haemophilus Influenzae Type b Combined	Intramuscular / Left thigh
	Hep B (3) Hepatitis B Vaccine	Intramuscular / Right thigh
	PCV ₇ (2) Pneumococcal Conjugated Vaccine	Intramuscular / Right thigh
6 MONTHS	Measles Vaccine (1)**	Subcutaneous / Left thigh
9 MONTHS	PCV ₇ (3) Pneumococcal Conjugated Vaccine	Intramuscular / Right thigh
12 MONTHS	Measles Vaccine (2)**	Subcutaneous / Right arm
18 MONTHS	DTaP-IPV // Hib (4) Diphtheria, Tetanus, Acellular Pertussis, Inactivated Polio Vaccine and Haemophilus Influenzae Type b Combined	Intramuscular / Left arm
6 YEARS (BOYS & GIRLS)	Td Vaccine Tetanus and Reduced Strength of Diphtheria Vaccine	Intramuscular / Left arm
9 YEARS (GIRLS ONLY)	HPV Vaccine	Intramuscular / Arm or thigh
12 YEARS (BOYS & GIRLS)	Td Vaccine Tetanus and Reduced Strength of Diphtheria Vaccine	Intramuscular / Left arm

*Rotavirus vaccine should NOT be administered after 24 weeks.

**Do not administer with any other vaccine.

DROPS BY MOUTH

LIQUID BY MOUTH

INJECTION

PROFMED

0860 776 363
profmedbaby@profmed.co.za